# MODELLING OF ANTIBIOTIC PRODUCTION BY *BACILLUS LICHENIFORMIS* USING A STRUCTURED/SEGREGATED APPROACH

APPROVED BY

DISSERTATION COMMITTEE:

Supervisor: ______________________

______________________

______________________

______________________

______________________

This dissertation was prepared with the LaTeX document preparation system using additional macros written by John W. Eaton based on the thesis style written by Khe-Sing The for The University of Texas at Austin.

TeX is a trademark of the American Mathematical Society.



Printed in the United States of America.

10 9 8 7 6 5 4 3 2 1

To Diana.
You make it all worthwhile.

# MODELLING OF ANTIBIOTIC PRODUCTION BY *BACILLUS LICHENIFORMIS* USING A STRUCTURED/SEGREGATED APPROACH

**by**

**ANDREW PATRICK FORDYCE, B.S. Ch.E.**

**DISSERTATION**

Presented to the Faculty of the Graduate School of

The University of Texas at Austin

in Partial Fulfillment

of the Requirements

for the Degree of

**DOCTOR OF PHILOSOPHY**

THE UNIVERSITY OF TEXAS AT AUSTIN

May1992

# Acknowledgments

I have received an immense amount of help and support during my time at The University of Texas. I would like to acknowledge those people who were most instrumental in my success.

Special gratitude is owed to my research advisor, Professor James B. Rawlings. Through his advice I was able to gain many insights concerning my project. I learned a great deal from observing his approach to solving engineering problems.

I would like to thank my dissertation committee members: Professors Thomas Edgar, James Brock, George Georgiou, and Alan Schauer. Special thanks to Professor Georgiou, who graciously allowed me complete freedom in using his laboratory.

Someone once told me I would learn more in graduate school from the students than the professors. Someone was right. I would like to thank W. Witkowski, S. Miller, T. Mullins, K. Muske, and H. Matthews for sharing some of their intelligence with me. John Eaton deserves special acknowledgment for keeping my head above the computational backwaters.

I have been fortunate enough to meet some good friends in Austin. Thanks to Allen Hoffman, Bill Wagner, and John Knoblauh for the hilarious times. Tony and Jodee Mullins deserve my sincere gratitude. Their unselfish generosity made my last months in Austin tolerable. I would also like to thank the Bytnar family for making me feel welcome in their home.

These acknowledgments would not be complete without mention of my family. The love and support my parents and siblings have given me throughout my life have inspired me to be my best.

I would like to thank my wife, Diana. Her unfailing love and belief in me convinced me I could succeed.

Finally, I thank God for staying with me throughout the difficult times.

Andrew Patrick Fordyce

*The University of Texas at Austin*
*May1992*

**MODELLING OF ANTIBIOTIC PRODUCTION BY *BACILLUS LICHENIFORMIS* USING A STRUCTURED/SEGREGATED APPROACH**

Publication No. ________________

Andrew Patrick Fordyce, Ph.D.
The University of Texas at Austin, 1992

Supervisor: James B. Rawlings

This study evaluates the effectiveness of a segregated model for prediction of differentiation-related secondary metabolism in a bacterial system. Specifically, this work has examined the production of the peptide antibiotic bacitracin by *Bacillus licheniformis* in a submerged-culture fermentation system. A segregated model recognizing the three morphological stages of the *Bacillus* life cycle has been developed. The sporangium biomass has been characterized using an age-population model to reflect the age-dependent antibiotic kinetics. Constitutive relationships governing the rates of vegetative cell reproduction, spore germination, commitment to sporulation, and substrate consumption have been proposed. Based on this model framework, the dynamic cell growth, differentiation, and antibiotic production equations have been developed. Numerical solution techniques have been designed to provide solutions to the steady-state and dynamic fermentation equations.

As a means of comparison between the segregated modelling approach and more common fermentation models, a structured model has been adapted to the bacitracin system. The comparison model utilizes a two-compartment age model to describe cell growth, differentiation and bacitracin production.

Batch, steady state, and step-test fermentation data from a laboratory-scale fermentor have been incorporated into a maximum likelihood parameter estimation scheme for model identification. Confident estimates of the growth and differentiation parameters have been obtained for the segregated model using available measurements. However, the insensitivity of the bacitracin data to the production function only allowed for confident determination of a constant antibiotic production coefficient.

In contrast to the segregated model, the structure of the comparison model prevented complete utilization of the available measurements in the parameter estimation scheme. This created a situation in which the information provided by the applicable measurements was insufficient for confident determination of the model parameters. The two-compartment age model was also unable to adequately describe the delay between fermentor inoculation and the appearance of antibiotic.

Open-loop optimal control studies were utilized to compare the suggested control policies of the two models. The computed strategies were significantly different, with the segregated model suggesting a non-batch feeding policy that predicts improved antibiotic yield of 30% over the comparison model batch policy.

# Table of Contents

# List of Tables

# List of Figures

# Chapter 1

# Introduction

Products derived from microbial sources have played an important role in the development of our society. A significant class of microbially-derived products are the secondary metabolites, also known as idiolite compounds. The secondary metabolites are considered to include all metabolic compounds that do not play a direct role in the normal activities of an organism. While the role of these compounds in the producing organisms is an interesting question that has stimulated much research, the secondary metabolites have achieved a higher level of importance due to fact that many of these compounds have found applications in human and animal medicine. An example of the significance of the secondary metabolites is the realization that most antibiotic compounds used in the treatment of bacterial infection are derived from compounds that are classified as idiolites.

The economic importance and increased demand for many secondary metabolites has stimulated the search for improved production strategies. Currently, these strategies have concentrated on screening for higher producing microorganisms, generation of more efficient microbial strains through mutagenesis, and improvement of growth and production media. An underutilized means by which bioprocesses can be improved is the application of process control strategies. In this context, the term control strategy is meant to imply manipulation of the available process operating variables such that some predetermined measure of process performance is improved. The basic reason that advanced control strategies are rarely utilized to improve bioprocess performance is because a high level of process understanding is essential to the success of the control scheme. Normally, process understanding is represented in the form of a mathematical model that a model-based control scheme can utilized to make decisions on how to manipulate operating variables to best achieve the desired operating goal. The complex nature of microorganisms makes them difficult systems to model. Nonlinear dynamic behavior, complex input interactions, and large time delays are common features of microbial systems.

Current efforts at developing advanced-control schemes for secondary metabolite systems have utilized simple unstructured microbial growth and product production models. These unstructured models assume microbes are a single component distributed continuously through the fermentation medium. This oversimplified picture of a fermentation system generally results in a poor characterization of observed secondary metabolite production behavior. This in turn degrades the ability of control algorithms to optimize process performance.

In actuality, microorganisms are individual entities consisting of a heterogeneous mixture of many different chemical species. Therefore, compared to the unstructured description, a more accurate picture of a fermentation system would be to recognize that a population of individual organisms exists, where the behavior of each individual cell contributes to the behavior of the population as a whole. Models that recognize the existence of individual cells and account for their effects on the cell populations are known as structured-segregated models.

Historically, the utility of population models has been limited due to several factors. A primary barrier is the lack of understanding of the mechanisms involved in cell growth and product production, making it difficult to formulate useful segregated model descriptions of fermentation systems. A second problem is the computationally-intensive nature of solution techniques required to solve the modelling equations resulting from the population balance. Further, the difficulty in obtaining measurements of cell population characteristics has been perceived as a limitation in the ability to identify segregated models. However, advances in molecular biology and the increased computation capabilities of digital computers have begun to remedy the basic problems with the segregated modelling approach.

Considering the desire to develop effective control strategies for secondary-metabolite fermentations, the goal of this work has been to develop a segregated model for secondary metabolite production and to analyze the advantages and disadvantages of the segregated approach compared to less sophisticated fermentation models. Due to the diverse nature of secondary metabolites and the organisms that produce them, it is difficult to make generalizations concerning the utility of a particular modelling approach for characterizing all secondary-metabolite fermentations. Therefore, to narrow the scope of this research to a manageable level, a specific fermentation system, representative of an industrially important subclass of secondary metabolites, will be the focus of the work presented in this thesis.

Specifically, this work develops a structured population model for the production of the peptide-antibiotic bacitracin from the bacterium *Bacillus licheniformis*. Included in this development is analysis of the issues involved in model identification, and an examination of the ability of the model to characterize fermentation behavior under a range of operating conditions. To provide a means of comparison between the proposed segregated model and the less sophisticated modelling approaches, a model from the fermentation literature has been adapted to the bacitracin system.

## 1.1 Review of Secondary Metabolite Production Models

Prior to a detailed development of the segregated growth and antibiotic production model, a review of the modelling approaches in the fermentation literature is helpful. Due to the economic importance of secondary metabolites, many researchers have attempted to model the growth and production characteristics of organisms that form secondary metabolites. Many different modelling approaches have been utilized to characterize secondary metabolite systems. The following discussion provides a review of the idiolite production models.

### 1.1.1 Unstructured Models

The simplest modelling approach, known as unstructured, is to approximate the microbial culture as a single chemical compound distributed evenly throughout the fermentation medium. Unstructured models do not recognize any measure of the culture viability, nor the fact that a population of individual microbes exists. The unstructured approach assumes that the system

| Model Type | Product | Reference |
|---|---|---|
| Luedeking and Piret | penicillin | Constantinides et. al. 1970 [19] |
| Monod | griseofulvin | Calam et. al. 1971 [15] |
| Luedeking and Piret | penicillin | Calam and Russel 1973 [16] |
| substrate inhibition | penicillin | Hegewald and Rückbeil 1978 [49] |
| Monod multisubstrate | penicillin | Giona et. al. 1976 [40] |
| Monod | exotoxin | Holmberg et. al. 1980 [53] |
| Monod multisubstrate | penicillin | Bajpai and Reuß 1980 [5] |
| Monod multisubstrate | oxytetracycline | Shnaider and Biryukov 1985 [95] |
| catabolite repression | $\alpha$-amylase | Yoo et. al. 1988 [113] |

**Table 1.1**: Examples of unstructured idiolite production models.

behaves independently of its history, and the culture can be characterized using a measure of the quantity of biophase, such as biomass concentration or dry weight [48]. Implicit in the unstructured model is the assumption that balanced growth prevails at all times. A growth period is called balanced if every extensive variable of the system increases at the same rate. With this assumption, a single measure of the culture, e.g., biomass concentration, is sufficient to characterize all changes during the growth period [34].

The simplicity of the unstructured model has led to its historical dominance in the fermentation modelling literature. Table 1.1 provides a sampling of the unstructured models applied to idiolite producing systems. The first attempt at utilizing an unstructured model for a secondary metabolite fermentation has been attributed to Luedeking and Piret [64]. This classic model assumes that the rate of product formation is proportional to a combination of biomass concentration and the biomass growth rate:

$$\frac{dP}{dt} = a_1 X + a_2 \frac{dX}{dt} \tag{1.1}$$

where $P$ is the product of interest, and $X$ is the biomass concentration. This function takes into account growth-related production along with non-growth associated enzymatic activity. The Luedeking and Piret model has been utilized to model penicillin production by Constantinides et al. [19] and Calam and Russel [16]. A similar model was used by Holmberg et. al. [53] to describe exotoxin production by *B. thuringiensis.* One shortcoming of this modelling approach is its inability to predict a time delay between culture inoculation and the onset of product formation. The groups using this approach for penicillin production assumed a 20 hour delay between culture inoculation and penicillin production. Holmberg et. al. estimated that approximately 5 hours delay occurred before exotoxin appeared in batch culture. A second problem of modelling product formation based only on growth rate and biomass concentration is evident in fed-batch culture where the addition of easily metabolizable sugars can lead to unmodelled catabolite repression of product formation.

In an attempt to model the catabolite repression effect, and to explicitly recognize the dependence of idiolite production on substrate concentration, several groups have utilized unstructured kinetic models. In this approach, product formation is a function of the culture conditions:

$$\frac{dP}{dt} = f(\mathbf{s}, X) \tag{1.2}$$

where $\mathbf{s}$ is the substrate concentration vector, and $f(\mathbf{s}, X)$ is a general non-linear function relating substrate and biomass concentration to the product formation rate. Calam et al. [16], Hegewald and Rückbeil [49], Yoo et. al. [113], and Shnaider and Biryukov [95] have considered a variety of limiting substrate kinetic models including Monod, catabolite repression, and substrate inhibition kinetics in attempting to model several different antibiotic and enzyme production fermentations. Bajpai and Reuß [5] and Giona et al. [40] both utilized models where the rate of penicillin formation depended on limiting sugar and dissolved oxygen concentration.

### 1.1.2 Structured Models

The structured modelling approaches have emerged as an attempt to provide a better measure of the culture state, thus allowing for dependence of current behavior on culture history, and allowing for situations where unbalanced growth prevails. Reviews of structured microbial growth models have been provided by Harder and Roels [48] and Shuler [98]. Structured models treat the culture not as a single entity, but as groups of distinguishable entities, which taken together represent the biotic phase. While these models can account for time-dependent changes in the makeup of the biotic phase, they cannot account for variation between individual organisms within the population. However, the structured models do provide a better approximation to the actual culture state then an unstructured approach. Chemical structure is the most common type of structured model. An example would be a model tracking the concentration of RNA and protein [87]. This two compartment chemical model is more complex than the unstructured model allowing for a measure of the biomass quality, rather than just a simple measurement of biomass quantity.

Several applications of structured models to secondary metabolite forma-

| Model Structure | Product | Reference |
|---|---|---|
| age | antibiotics | Aiba and Hara 1965 [1] |
| morphological | antibiotics | Megee et. al. 1970 [72] |
| age | gramicidin-s | Blanch and Rogers 1971 [11] |
| age | streptomycin | Brown and Vass 1973 [13] |
| chemical | exoenzymes | Van Dedem et. al. 1973 [24] |
| age | antibiotics | Fishman and Biryukov 1974 [31] |
| age | macrolides | Martin and McDaniel 1975 [69] |
| morphological | cephalosporin C | Matsumura et. al. 1981 [71] |
| chemical | alkaloids | Patžoutova et. al. 1981 [80] |
| morphological | $\delta$ endotoxin | Schulz et. al. 1985 [93] |
| chemical | tetracycline | Běhal et. al. 1985 [14] |
| chemical | cycloheximide | Dykstra et. al. 1988 [27] |

**Table 1.2**: Examples of structured secondary metabolite production models.

tion, shown in Table 1.2, have appeared in the literature to date. Examples of chemically structured models are common. One example is the production of alkaloids by *Claviceps purpurea*. Pažoutova et al. [80] utilized a chemically structured model that accounts for the intracellular phosphate concentration as an indicator of the product formation rate. Other researchers have used catabolite repression models to track the intracellular concentration of some critical product synthetase enzyme. Dykstra et al. [27] used this approach for cycloheximide production by *Streptomyces griseus.* Běhal et al. [14] modelled tetracycline production by predicting the activity of an oxygenase enzyme, considered limiting in the tetracycline fermentation. Van Dedem and Moo-Young [24] have compared induction and repression mechanisms in the formation of extracellular enzymes using a structured model.

A second form of structured model is the age-structured model. A number of natural aging phenomena have been observed in microbial systems. Growth in many mycelial organisms occurs in the apical region. Bud scars in yeast cells establish a physical age marker that affects the ability of the yeast cell to reproduce. Cell differentiation has been attributed to an aging process and has been linked to product formation [96]. In a general formulation, a model based on age would be considered segregated because age is a continuous variable providing a unique identity to each cell in the culture. Several authors have reduced the general age-segregated model to a structured conceptualization by assuming the continuous age distribution can be approximated by two or more age groups. Blanch and Rogers [11] used an age-structured model to predict gramicidin S formation by *Bacillus brevis.* They divided the cell culture into two groups, mature and immature cells. Mature cells were responsible for both antibiotic formation and reproduction. This model was able to simulate experimental fermentation responses for both batch and continuous culture, except at high dilution rates. The authors suggest several possible explanations for the poor performance at high dilution rate, including unmodelled growth lags, and chemostat instability. Brown and Vass [13] used a similar age-structured model for gramicidin S and streptomycin fermentations. Martin and McDaniel [69] also used the age grouping technique to model macrolide antibiotic production by a variety of *Streptomyces* species.

A minor variation of the age-structured model, introduced by Aiba and

Hara [1], correlates the culture's "cumulative age" to the specific antibiotic formation rate. The cumulative age is a crude measure of the average age of the cell population. The cumulative age, $\theta$, was given by:

$$\theta = \frac{X_0\theta_0 + \int_{t_0}^{t} \tau X(\tau)\, d\tau}{X(t)} \tag{1.3}$$

where $X_0$ and $\theta_0$ are the initial cell concentration and cumulative age, respectively. Fishman and Biryukov [31] used this approach fitting a quadratic function of cumulative age to the specific product formation rate and developed feeding strategies for maximizing product.

Models that can be considered structured, but do not use chemical or age grouping have also appeared in the literature. Megee et al. [72] designed an idiolite production model based on the different mold morphologies observed in submerged culture. This model takes advantage of the common observation that secondary metabolites appear concomitant with cellular differentiation processes. A similar approach was used by Matsumura et al. [71] to characterize the production of cephalosporin C by *Cephalosporium acremonium*. This group divided the biotic phase into three separate morphological groups: hyphae, swollen fragments, and arthrospores. From experimental observation it was noted that the swollen fragments were the highly productive fraction of the biomass. Using the morphologically-structured model they were able to closely approximate experimental antibiotic production data. A final morphologically-structured example is provided by Schulz et. al. [93] who used several morphological states to model the $\delta$ endotoxin formation in spores of *B. thuringiensis*.

### 1.1.3 Segregated Models

A more accurate portrait of a microbial system is a population of cells, each cell having its own distinguishing characteristics, each cell responding to its local environment. The segregated modelling approach attempts to characterize the fermentation based on the idea that the overall population behavior can be attributed to the behavior of the individual members of the population. This methodology can be used to account for phenomena such as cell-size dependent growth rates, age-dependent production rates, or any other activity which can be related to some characteristic that distinguishes members of the population. The segregated modelling approach, known as population modelling, has been reviewed by Ramkrishna [86, 85] and a general discussion is provided by Hulburt and Katz [54].

Very few population models have been applied to secondary metabolite fermentations. One possible reason for the dearth of population models for secondary metabolite systems is the difficulty in measuring distributions of population characteristics in fermentation systems [97, 5]. One exception is the model presented by Shu [96]. Shu used an age-segregated population balance to model product accumulation in batch and continuous fermentation systems. The specific product formation rate was fit using a set of exponential basis functions, given by:

$$\mu_p = \sum_i A_i \exp(-k_i \tau) \tag{1.4}$$

where $\tau$ is the cell age and $A_i$ are experimentally fit parameters. This formulation is fairly versatile, and it can be used to model the transient behavior of many types of fermentations. Some drawbacks of this approach include difficulty in estimating the parameters of the production function and the fact that this model does little to explain the physical mechanisms involved in secondary metabolite formation [97].

## 1.2 Research Goals

At this time, there are few experimental studies concerning the application of a population balance model to a microbial system. It is the objective of this research to help fill this void by examining the feasibility and utility of using a segregated model to predict the dynamic behavior of a microbial secondary metabolite fermentation system. As a part of this project, the following research goals were identified:

- To identify a secondary metabolite system for which a segregated model could facilitate improved predictive capabilities.
- To develop a dynamic model describing the growth and product production using a segregated model.
- To construct a laboratory–scale fermentation apparatus to collect data for model identification.
- To examine the accuracy in different regions of the operating space.
- To compare the proposed model to a structured model existing in the fermentation literature.
- To analyze and compare open-loop optimal control strategies for maximization of bacitracin production based on each model.

## 1.3 Dissertation Organization

This dissertation is divided into six chapters. This chapter contains introductory material concerning the author's perception of the role of mathematical modelling in the realm of fermentation engineering. This includes a literature review of fermentation models for secondary metabolite production, and a discussion of the goals of this research project.

Chapter 2 proposes modelling of the production of the peptide antibiotic bacitracin by *Bacillus licheniformis* as a model system to explore the utility of the segregated modelling approach. Based on the biology and physics of the proposed model system, a dynamic, segregated model is developed, and solution techniques for the resulting model are described. To provide a comparison for the segregated approach, a structured model proposed by Blanch and Rogers [11], is modified to describe the bacitracin system.

The third chapter describes the experimental apparatus used for model identification and validation. A detailed description of the laboratory-scale fermentor is given, along with discussion of the experimental procedures for growth and analysis of the bacteria and the product assays.

Chapter 4 examines the issues of model identification and validation. Model parameters for growth, differentiation, and antibiotic production are estimated for both the segregated and comparison models, based on experimental fermentation data. Evolution of the models from their initial proposed forms to their final form is discussed. The resulting parameter estimates are examined.

In Chapter 5 comparisons of the two models are made. Included in the comparison are issues of model identification, effectiveness of model fit, and open-loop optimal control strategies suggested by each model.

The final chapter summarizes the conclusions of this research, and suggests directions for further study.

# Chapter 2

# Model Development and Solution

## 2.1 Model System

The basic goal of this research is to explore the effectiveness of the segregated modelling approach for the prediction of secondary metabolite production. In selecting a model system several criteria were considered important. First, to analyze the utility of a segregated model it was necessary that suitable non-segregated models existed in the literature to provide a yardstick for comparison purposes. Additionally, since experimental verification of a model is desirable, the growth of the model organism and metabolite production needed to be experimentally tractable. Finally, it was important that the chosen metabolite product be industrially relevant. With those considerations in mind, production of the peptide antibiotic bacitracin by the bacteria *Bacillus licheniformis* was chosen to explore the effectiveness of the segregated modelling approach.

Bacitracin production is a good choice as an experimental model for many

| Product | Application |
|---|---|
| peptide antibiotics | feed additives, Topical Antibiotics |
| amino acids | nutritional supplements |
| exoenzymes | food, textiles, paper Industries |
| insect pathogens | biopesticides |
| biosurfactants | enhanced oil recovery |

**Table 2.1**: Industrial applications of sporulation-associated products from *Bacilli* [23, 35].

reasons. First, many of the useful *Bacillus* products are associated with the differentiation process, and insight into strategies for production of bacitracin could be directly applicable to a wide variety of *Bacillus* derived metabolites, including the sporulation-related products shown in Table 2.1. Concerning experimental tractability, species from the *Bacillus* genus generally have simple nutritional needs and grow quickly. A large literature exists concerning growth conditions, and procedures for the assay of bacitracin are available. Also, the *Bacillus* unicellular morphology greatly simplifies the measurement of cell numbers. Finally, although no structured models have appeared for the production of bacitracin, an age-structured model, conceived by Blanch and Rogers [11], exists for the production of a closely related peptide antibiotic, gramicidin. This model is adaptable to the bacitracin system, and serves as a comparison model for the segregated approach.

The genus *Bacillus* includes the aerobic, unicellular endospore forming bacteria. These microorganisms are normally gram positive, but some species become gram negative upon differentiation. All species can utilize oxygen in




**Figure 2.1**: The *Bacillus* life cycle consists of three stages. The first is the vegetative stage, in which cell growth and fission occur. In nutrient depleted surroundings sporangium cells form, starting the cell differentiation process. A mature endospore is the final stage of the cycle.

aerobic respiration, while some species can also take advantage of anaerobic fermentative and denitrificative pathways [103]. The *Bacillus* life cycle, illustrated in Figure 2.1, begins with germination of an endospore cell when environmental conditions become favorable. The resulting vegetative cell reproduces via binary fission until the environment becomes adverse to normal growth. At such a time, the cell begins a differentiation process converting from a vegetative cell to a sporangium. In a sequence of well-timed steps a single endospore forms within the sporangium cell. Concurrent with the differentiation process many secondary metabolites, including several types of antibiotics and exocellular enzymes, are produced. When the endospore reaches maturity the sporangium cell lyses, releasing a mature spore. The spore remains in a state of cryptobiosis, with no detectable metabolic activity, until environmental conditions become favorable for growth again [103].

### 2.1.1 Review of Sporulation Models

Considering the connection between idiolite production and the differentiation process exhibited by *B. licheniformis*, a prerequisite of predicting antibiotic production is the ability to model the dynamics of growth and differentiation. At this time, only a few mathematical models for the *Bacillus* growth and differentiation processes have appeared. The first is attributable to Dawes and Thornley [22], who developed a segregated model to describe growth and sporulation of *B. subtilis* in continuous culture. More recently, sporulation models have been examined by Ollis [79] and Schulz et. al. [93]. Ollis modelled the production of exotoxin from *B. thuringiensis* using an age-structured model. The kinetics of exotoxin production were treated using the Luedeking-Piret equation (see Section 1.1.1), and spore formation was considered to be first order with respect to the delayed biomass concentration. Ollis' model results in 3 ordinary differential equations coupled to a delay differential equation for the spore concentration. The age-structured model was able to model batch data well. However, the simple spore initiation model used could lead to problems in continuous and fed batch fermentations, where the addition of easily metabolizable sugars leads to repression of sporulation commitment.

Similar to Ollis, Schulz et. al [93] modelled exotoxin production from *B. thuringiensis* using a morphologically-structured model with four distinguishable cell states. Transition between the vegetative phase to sporangium, and in turn, to endotoxin containing cells, and finally mature spores was modelled using delay differential equations. This model was used to characterize batch and continuous growth, and optimal operation was determined for one and two-stage continuous fermentations.

The development of highly-structured, single-cell models, popularized by the Cornell *E. coli* model of Shuler et al. [99], have recently been applied to *Bacillus* species in an attempt to model the differentiation process. Jeong et. al. [56] have proposed a structured single-cell model consisting of 39 nonlinear differential equations with almost 200 parameters. The goal of this model is to account for the principal interactions between the major metabolic networks in *B. subtilis*, with the goal of describing the biochemical patterns associated with the transition from vegetative growth to sporulation. A similar structured approach has been used by Starzak and Bajpai [104] to model the vegetative growth cycle and the transition to sporulation. Neither of the highly-structured models make a clear distinction between cells in the vegetative or sporangium stages of development, instead commitment to sporulation is treated as a continuous transition with no clear demarcation between growth and differentiation The inability of these models to predict simultaneous presence of vegetative and differentiated cells is an obvious shortcoming, but the basic structure of these models does allow for an in depth analysis of the intracellular dynamics of growth and the control system for sporulation initiation.

## 2.2 Segregated Model Development

In contrast to the goals of the single-cell models proposed by Jeong et. al. [56] and Starzak and Bajpai [104], the intended use of the model proposed in this research is as a tool for development of control strategies. Considering this, it was determined desirable properties of the model included a minimum of adjustable parameters and reasonable computational times, while still main-




**Figure 2.2**: A well-mixed, stirred-tank fermentor depicting a microscopic view of a population of *Bacillus licheniformis*.

taining an accurate representation of the physical system.

### 2.2.1 Microscopic View

A typical *Bacillus* fermentation consists of a mixture of cells in an aqueous environment containing a variety of soluble nutrient components and metabolic byproducts. A view of a microscopic fluid packet, depicted in Figure 2.2, shows a collection of cells in various stages of growth and differentiation. Each cell consists of many different chemical components with a multitude of simultaneous reactions, allowing the cell to both affect and respond to its local environment. The situation is further complicated by a variety of mass transfer processes involved in the internalization, transformation, and incorporation of nutrients into existing and new biomass. Based on this description, it is easy to imagine how overwhelming a complete mechanistic model of all reactions and transport processes would be. Therefore, the task is to determine the essential elements of the fermentation, and to translate those processes into a mathematical description. In this way, the complexity of the problem is reduced, and the important aspects of the system are captured.

In an effort to develop a useful, yet tractable, model for production of bacitracin from *B. licheniformis*, inspiration is taken from the microscopic picture of a culture. An obvious characteristic of a *Bacillus* culture is the presence of three distinct cell types: vegetative, sporangium, and spore, as shown in Figure 2.1. A logical modelling approach is to structure the biomass according to its three distinct morphologies. Morphological structure is appealing because it provides an accurate representation of the actual microscopic picture. Also, since bacitracin production is localized in the differentiating portion of the biomass, it is important that a model recognize the existence of the different types of biomass. Dawes and Thornley [22] have utilized morphological structure to develop a steady-state model for *Bacillus subtilis* in continuous culture. In this work, a similar morphological structure has been utilized to develop a dynamic model for *Bacillus licheniformis* in batch and continuous culture modes.

### 2.2.2 Spore Biomass

Species from the *Bacillus* genus are ubiquitous in soil environments and exist primarily in a dormant state, known as a spore. The spore morphology is ellipsoidal and approximately 0.3–0.5 $\mu$m in diameter. This is distinctly different than the rod-shaped vegetative cell morphology. Using phase-contrast microscopy spores appear as phase-bright bodies. In comparison, the vegetative cell form appears as dark, elongated body, and thus the presence of spores is readily detectable using light microscopy.

In addition to the visible morphological differences, spores also exhibit significantly altered resilience with respect to harsh environmental conditions. This includes an abnormally high resistance to heat, desiccation, radiation, and chemicals. The unusual resistance properties of spores are attributable to several major differences between the outer layers of a spore and those of vegetative cells. Normal gram-positive vegetative cells are contained by a phospholipid bilayer enclosed by a rigid peptidoglycan cell wall. In addition to a cell wall and membrane, spores are enclosed by several extra layers, including the cortex, spore coat, and exosporangium. It appears that these additional layers, along with several stabilizing adaptations of the spore protoplast, are responsible for the acquisition and maintenance of resistance properties, allowing for sustained viability in unfavorable environmental conditions [38].

The spores remain protected in a dormant state, until environmental conditions change such that growth becomes favorable. At this time, the spore sheds its protective layers and returns to the vegetative state.

### 2.2.3 Vegetative Biomass

In a normal growth situation a *B. licheniformis* culture consists primarily of vegetative cells. These cells represent the portion of the biomass that is actively growing and dividing. Vegetative cells are gram positive and are rod shaped, measuring approximately 0.6 $\mu$m in diameter and 2.2 $\mu$m in length. The mode of reproduction is binary fission, resulting in the production of two daughter cells of approximately equal size. Vegetative cells can exist in short chains of 2–3 cells, however, in a well-mixed environment the majority of the cells exist as singlets. Most strains exhibit motility, particularly in nutrient depleted conditions [41].

*B. licheniformis* is a facultative aerobe, having a functional respiratory chain, allowing for growth using electron transport with oxygen as the terminal electron acceptor. In anaerobic conditions, *B. licheniformis* utilizes either an anaerobic respiratory pathway, with nitrate as the electron acceptor, or a fermentative pathway yielding a mixture of organic acids [59]. Vegetative growth is possible using a variety of carbon and nitrogen sources in defined and complex media, and typical doubling times range from 50–90 minutes, depending on the nutrient source and oxygen availability [59].

### 2.2.4 Sporangium Development

In addition to vegetative cells and spores, a third type of cell exists in a pure *B. licheniformis* culture. These cells, known as sporangia, represent the individuals in transition between the vegetative growth state and the dormant spore.

After commitment to the sporulation pathway, the cell goes through a tightly regulated differentiation process coded in the DNA. Historically, the progress of sporulation has been gauged based on morphological changes visible by microscopy. The morphological markers of sporulation are depicted in Figure 2.3. Stage 0 is the vegetative state, and therefore, represents a growing cell, prior to commitment. There exists controversy concerning the first morphological manifestation of sporulation. Prior to mutational studies, stage I was consider to be condensation of the chromosome in an axial filament. However, since no mutants have been found to be deficient in formation of the axial chromosomal filament, stage I has been generally discarded [63]. The first accepted morphological development, termed stage II is the appearance of an asymmetric membrane division, dividing the cell into two compartments of unequal size. Both the small compartment, known as the forespore, and the large compartment, the mother cell, contain intact copies of the chromosomal DNA. At this point, the sporulation sequence progresses to stage III, where the small compartment is engulfed by the mother cell, producing a free protoplast within the mother cell's cytoplasm. The protoplast has a double-walled membrane created during engulfment. During stage IV a thick layer of modified-peptidoglycan wall material is deposited between the two forespore membranes, creating the spore cortex. Stage V brings enclosure of the outer membrane layer in a covering of polypeptides, called the spore coat. At this time, the spore develops its characteristic refractility, and becomes visible using phase-contrast light microscopy. Following the development of refractility, the prespore develops into a mature spore, gaining the resistance properties against heat, chemicals, desiccation and radiation. The stage of spore maturity is termed stage VI. The final development, stage VII is lysis of the mother cell releasing a mature spore. The entire sporulation process, from commitment to appearance of mature free spores takes approximately 8–10 hours [76, 63].

Along with the characteristic morphological changes observed during differentiation, a variety of biochemical markers are also recognized in association with stages of spore formation. Production of extracellular macromolecular degradation enzymes including $\alpha$–amylase, proteases, and nucleases are all associated with early sporulation (i.e. stages 0–II). Antibiotics, including bacitracin in *B. licheniformis*, are also expressed early in the differentiation sequence [76]. Mid-sporulation biochemical markers include alkaline phosphatase at stage II and glucose dehydrogenase between stages III and IV. A pronounced late-sporulation marker is the expression of dipocolinic acid, which is unique to spores [83].




**Figure 2.3**: The morphological markers of spore formation.

### 2.2.5 Model Structure

By deciding to structure the *B. licheniformis*-bacitracin system based on cell morphology, a foundation for the proposed model has been established. In addition to the model structure, it is necessary to characterize the level of sophistication required to model each portion of the biophase. Finally, to complete the model, it is essential that the mechanisms by which the cell groups interact with each other and their environment are identified and characterized.

The interconnect cycles of growth and differentiation exhibited by *B. licheniformis* are diagrammed in Figure 2.4. Processes including spore germination,




**Figure 2.4**: The interconnected cycles of growth and differentiation in *B. licheniformis*. Cycle 1 depicts the process of vegetative reproduction and its connection to the consumption of substrate, $s$. Cycle 2 represents the differentiation cycle, and its connection to the production of antibiotic, $A$. The rates of spore germination, vegetative growth, sporulation commitment, and antibiotic production are represented by $\mu_g$, $\mu_v$, $\mu_s$, and $\mu_a$, respectively.

vegetative growth, substrate consumption, and sporulation commitment are all involved in the relationships relating cell morphologies, and therefore must be described by the model. Further, the relationship between the process of differentiation and production of bacitracin is essential in attaining the desired goal of modelling bacitracin production.

In the following sections, a model for each of the three portions of the biomass is developed. Also, mathematical representations for the growth and differentiation rate processes are examined. The combination of the rate expressions and the structure of the biophase comprise the kinetic portion of the segregated mathematical model for the *B. licheniformis* fermentation system.

### 2.2.6 Vegetative Biomass Model

In its original formulation the Dawes and Thornley model [22] was primarily for description of steady-state growth and sporulation kinetics found in the chemostat fermentor. Therefore, a Monod type vegetative growth law was used to represent the balanced growth conditions of the chemostat. Assuming growth of vegetative cells, $X_v$, is limited by a single substrate, of concentration $s$, then the Monod growth law is of the form:

$$\frac{dX_v}{dt} = \mu_v X_v \tag{2.1}$$

$$\mu_v = \frac{\alpha_v s}{K_v + s} \tag{2.2}$$

The Monod growth function, illustrated in Figure 2.5, models the specific rate of vegetative cell production using two parameters. The parameter $\alpha_v$ represents the maximum attainable growth rate, and $K_v$ is the substrate concentration that will support a growth rate of half the maximum. The structure of the Monod model is based on the assumption that growth is limited by a single biochemical pathway that exhibits Michaelis-Menten saturation kinetics. In most cases, this is an oversimplification of the actual situation, so it is not possible to link any phenomenological meaning to the growth parameters. However, the Monod expression has been successfully applied in many growth situations [4]. Utilization of Monod growth kinetics implicitly assumes vegetative cells are indistinguishable from each other, and the growth properties of the culture are based on an average cell. Further, balanced growth is a basic assumption of this approach. Based on the assumptions of indistinguishability and the lack of structure in the vegetative biomass, the Dawes and Thornley vegetative model is both nonsegregated and unstructured.

While this unstructured model of vegetative cells proved adequate under the steady-state conditions presented by Dawes and Thornley, batch growth




**Figure 2.5**: Monod saturation kinetics. The parameter $\alpha_v$ represents the maximum growth rate, and $K_v$ is the Michaelis constant representing the substrate concentration needed to sustain a growth rate of $\alpha_v/2$.

experiments carried out in this study showed the unstructured growth model was unable to describe many characteristics of transient batch growth. A detailed discussion of the experimental work used for parameter estimation is provided in Chapter 4. However, to provide a basis for the model development discussion, dimensionless cell number and substrate concentration data from a typical *Bacillus licheniformis* batch fermentation are shown in Figure 2.6. An obvious oversight of the pure Monod growth law is it inability to predict a decline in nonspore biomass associated with cell death. The batch data show a rapid decrease in the nonspore biomass without an associated increase in the spore count, thus implying not all vegetative cells form spores. This




**Figure 2.6**: Batch growth and sporulation data for *Bacillus licheniformis* showing transient growth phenomenon unexplained by a Monod growth law.

phenomenon is unaccounted for using Monod kinetics. A second more subtle inconsistency between the batch data and the unstructured growth law is the continued growth of the non-spore biomass after the limiting substrate has been exhausted. Finally, the initial phase of the fermentation, where little or no growth is apparent even though substrate is consumed, is also unexplained by an unstructured growth model.

**Structured Vegetative Growth**. While the morphological structure of the Dawes and Thornley model is still considered appealing, a more sophisticated vegetative growth model is necessary to adequately describe the transient




**Figure 2.7**: Structured vegetative cell model. First, external substrate, $s_e$, is internalized at a rate $\mu_i$, and converted to readily utilizable biosynthetic precursors, $s_i$. Second, cell precursor, $s_i$, is converted into new biomass, $X_v$, at a rate $\mu_v$.

growth phenomenon observed in the batch experiments. A two-compartment chemical model, diagrammed in Figure 2.7, has been proposed to describe the transient growth kinetics. The structured growth model makes a distinction between the ability of a cell to take up a limiting nutrient, and the capacity to convert that nutrient into biomass. The hypothesis of competitive rates of nutrient transport and biomass production was first proposed by von Bertalanffy (see [106] pg. 195). Similar two-compartment models have also been proposed by Williams [111] and Ramkrishna et. al. [87].

Using the structured vegetative model, growth is considered to be a two-stage process. First, the limiting nutrient is taken up by the existing biomass at a rate given by $\mu_i$. This nutrient exists internally in the cell as readily available precursors for biosynthesis and energy production. In the second stage, the internalized nutrient pool is utilized at a rate given by $\mu_v$, to produce biomass and energy necessary for growth and maintenance. The buildup of internal substrate, $s_i$, can be interpreted as an increase in average cell mass, and conversely a decrease in the internal metabolite pool occurs when the rate of cell division exceeds the rate of replenishment of biosynthetic capacity leading to a decrease in average cell mass. This concept is supported by the observation that cells growing in nutrient rich environments are large and become smaller when the surroundings become nutrient depleted.

The two-stage growth process provides a possible explanation for the transient growth phenomenon observed in the batch fermentation experiments. According to the structured vegetative model, the initial growth lag results because the newly inoculated cells must build biosynthetic capacity to utilize the available substrate. During the lag phase, the process of substrate internalization is the growth limiting step. A transition between the slow-growth lag period and the logarithmic growth phase occurs when enough substrate has been internalized to promote rapid growth. At that time, the rate limiting process shifts from substrate internalization to biosynthesis. The logarithmic phase continues until nutrient depletion forces a decline in the growth rate.

The two-compartment growth model also provides a possible explanation for post-nutrient exhaustion cell division observed in the batch fermentations. According to an unstructured vegetative growth model, cell growth rate immediately responds to changes in the external environment. This would suggest that cell growth would cease immediately upon exhaustion of the limiting substrate in the surrounding environment. Observation of batch fermentations shows that production of colony forming units continues for a short time after the limiting carbon source has been depleted in the growth medium. With the structured model, growth is dependent on the internal substrate concentration, thus allowing for situations where growth can continue in lieu of completely nutrient deficient surroundings. After the initial lag phase, the batch growth is limited by biosynthetic capacity. When the limiting substrate becomes exhausted in the growth medium, the internal biosynthetic capacity begins to decline. Thus, the rate-limiting process reverts back to the substrate internalization step, and growth slows until finally ceasing with the exhaustion of the internal nutrient supply. Therefore, the process of vegetative reproduction continues for a short period after the external limiting substrate is gone.

The death-associated decline in nonspore biomass can be modelled by including an death rate constant, $m_v$ in the biosynthetic grow rate function, $\mu_v$.

$$\mu_v(s_i) = \mu_t(s_i) - m_v \tag{2.3}$$

The explanation for this is that the total biosythetic rate, $\mu_t$, is a combination of growth and maintenance activity. Utilization of substrate for growth activity results in an increase in biomass. However, substrate is also consumed for energy and precursor generation necessary to retain viability. Therefore, substrate consumption resulting in increased biomass is only part of the total substrate consumption. Typically, the maintenance activity is considered to be constant, and thus results in a downward shift of the substrate dependent specific growth rate. The Monod growth law including a death term is shown in Figure 2.8. As the plot shows there exists a critical substrate concentration, below which the growth rate is negative. The negative growth rate represents loss of colony forming units resulting from cell death [4].

Utilizing the structured growth kinetics including death kinetics, the substrate internalization and cell growth rate expressions are given by:

$$\mu_i = \frac{\alpha_i s_e}{K_i + s_e} \tag{2.4}$$

$$\mu_v = \frac{\alpha_v s_i}{K_v + s_i} - m_v \tag{2.5}$$

The resulting equations utilize five adjustable parameters. Again, $\alpha_v$ and $\alpha_i$ represent the maximum rates of the respective processes and $K_v$ and $K_i$ are affinity constants representing the concentration needed for half the maximum rate. Cell death is accounted for by the endogenous metabolism term, $m_v$.

An additional consideration in modelling cell growth is the effect of the




**Figure 2.8**: Monod growth rate including death kinetics. The substrate concentration, $s_c$, is the critical substrate concentration below which cell death occurs.

antimicrobial product on the producing organism. In the case of *B. licheniformis*, Haavik [45] observed a minimal effect of bacitracin on growth rate for bacitracin concentrations up to 30 (IU/ml). In this study, maximal bacitracin titers reached approximately 3.0 (IU/ml), and therefore, the effect of bacitracin on growth rate was assumed to be negligible.

### 2.2.7 Spore Biomass Model

The spore biomass is a source for vegetative cells through the process of germination. Therefore, to provide a complete characterization of the *Bacil-*

*lus licheniformis* system, it is necessary to develop a rate expression relating germination to the fermentor conditions. The approach taken in this research is to treat the spores as indistinguishable and unstructured. In line with this simple conceptualization for the spore biophase, a Monod growth expression has been used to relate the specific rate of germination to the current limiting-substrate concentration:

$$\mu_g = \frac{\alpha_g s_e}{K_g + s_e} \tag{2.6}$$

As in the case of vegetative growth, $\alpha_g$ and $K_g$ represent the maximum germination rate and the germination Michaelis constant, respectively. Again, the biochemical significance of parameters $\alpha_g$ and $K_g$ is not obvious and this relationship is considered empirical. However, for most industrial fermentations, spore germination does not contribute significantly to vegetative cell concentrations after the initial inoculant is prepared [22]. Also, spore germination experiments, described in Section 4.3.1, indicate that germination is subject to a significant time delay. This observation is consistent with the findings of other groups. In particular, Maraheil et. al. [68] found that outgrowth in *B. brevis* occurred after a 8–10 hour lag period. Also, Dawes and Thornley [22] considered the number of germinating spores to be insignificant in continuous culture experiments with *B. subtilis.* These results indicate that germination is a relatively slow process compared to growth and sporulation in normal batch and continuous culture. Therefore, since the spores generally appear at the end of batch runs, when nutrient depletion has occurred, and there is a significant lag time during germination, it seems likely that spores contribute an insignificant number of cells to the vegetative phase. For this reason, little emphasis has been placed on the spore germination model.

### 2.2.8 Sporangium Model

As was discussed Section 2.2.4 on sporangium development, there are many morphological and biochemical differences between sporangia at different stages of development. Significantly, bacitracin is produced during sporulation, and the rate of bacitracin production for an individual sporangium is thought to depend on its progress through the sporulation pathway. Therefore, not only are the sporangia distinguishable based on their progress toward creating a spore, but it is logical that the model should recognize the differences in sporangia if the goal is to predict production of a sporulation dependent product, i.e. bacitracin. This implies the need for a segregated model, which recognizes the differences in sporangia, and allows for incorporation of the distinguishing characteristics of the sporangium population into the antibiotic production model.

**Population Balance.** Translation of the notion of distinguishability between individuals into a mathematical description involves development of the population balance. The basis of the mathematical description for a population is the definition of the particle density function, $f(\mathbf{z}, t)$. This scalar valued function is defined as the number of individuals per unit volume. The particle density expression is a function of $\mathbf{z}$ and $t$, where $\mathbf{z}$ defines a vector of distinguishing characteristics, and $t$ represents time. It is convenient to break the coordinate vector, $\mathbf{z}$, into two vectors, $\mathbf{x}$, representing the external or spatial coordinates, and $\mathbf{y}$, representing internal or descriptive coordinates. The need to include spatial coordinates comes from the fact that an individual could behave differently than others and is distinguishable due simply to its location in space. The internal coordinates represent the characteristics of an individual which make it distinguishable regardless of position. Examples of internal coordinates are size, age, and chemical composition. The choice of the internal coordinates allows the modeller infinite flexibility, but with increased coordinates comes increased complexity. Therefore, internal coordinates are chosen such that they reflect the most important characteristics of the system of interest.

After selection of suitable descriptive coordinates, the derivation of the population balance equations is simply an accounting of individuals with respect to the coordinate system of choice. General derivations of the microscopic population balance are provided by several researchers including Randolph and Larson [88], Rawlings [89], and Hulburt and Katz [54].

The population balance resulting from microscopic continuity is given by:

$$\frac{\partial f}{\partial t} + \nabla \cdot (f \frac{d\mathbf{z}}{dt}) = B - D \tag{2.7}$$

This equation describes the changes in the particle density function due to the changes in the coordinates, $\mathbf{z}$, and due to particles entering and leaving the population through birth and death mechanisms, $B$ and $D$ respectively.

**Age Population Model**. With the purported connection between differentiation and expression of bacitracin, a logical characterization of the sporangia population is some measure of progress through the differentiation process. As research into the sporulation process has progressed, it is becoming clear that differentiation in *Bacillus* is a precise sequence of temporally, spatially, and morphologically regulated events [62]. It appears that the steps of the differentiation process are controlled at the transcriptional level, with each set of genes in the sequence being activated at a specific time in the process [61]. This strict regulatory pattern forces a close correlation between the biochemical and morphological markers of sporulation and time since commitment to differentiation [63, 114, 91]. Therefore, a convenient means to characterize the sporangium population, based on stage of spore development, is the sporangium age, or time since commitment to sporulation.

Let $\tau$ represent the internal coordinate of sporangium age. Defining $X_s(\mathbf{x}, \tau, t)$ as the sporangium age density function allows for a segregated description of the sporangium population. Now, the remaining tasks to complete the population balance description is elucidation of birth and death mechanisms, by which cells enter and leave the population, and a description of how the population changes due to the fact that the internal coordinate, age, changes in time. Selection of cell age as an internal coordinate makes the latter of these two tasks a trivial matter. Cell age is directly correlated with time since birth, therefore, the rate that cells age, $d\tau/dt$, is a constant of one.

The choice of cell age as the descriptive coordinate in the population balance provides a convenient simplification to the modelling of sporangium birth. Since by definition, cells are of age zero at birth, all entries to the population resulting from birth mechanisms are localized at the zero age boundary. Therefore, an understanding of the rate that vegetative cells sporulate is sufficient information to describe the effect of birth on the sporangium population.

**Sporulation Commitment**. The mechanism by which vegetative cells begin differentiation and enter the sporangia age distribution is through sporulation commitment. Due to the utilization of bacterial sporulation as a paradigm for cellular differentiation mechanisms, great progress in the understanding of the genetics and biochemistry of sporulation has been made. A majority of the sporulation research to date has been done using a *Bacillus subtilis* experimental model. Assuming that *B. subtilis* and *B. licheniformis* have similar regulatory mechanisms for sporulation, a brief discussion of the current knowledge concerning sporulation is provided to help motivate the formulation of sporangium birth and maturation models.

In response to nutrient depletion the *Bacillus* have evolved several regulatory responses to attempt to adapt to the changes in the environment. These include the stringent response, induction of transport systems for alternative nutrient sources, expression of chemotaxis and motility mechanisms, secretion of macromolecular degrading enzymes, and induction of competence for transformation [102, 100]. These regulatory adaptations all represent ways the cell could possibly obtain an alternative nutrient source to replace the current dwindling supply. In the event that all efforts fail, the cell has a final avenue to preserve itself, that being activation of the sporulation pathway. By proceeding through a set of genetically predetermined steps, the vegetative cell transforms itself into a highly resistant storage vessel, containing the bacterial DNA and dormant metabolic machinery. Thus, by transforming into a spore, the cell effectively saves itself for a better day.

Although many aspects of sporulation remain a mystery, the basic regulatory mechanism for *Bacillus* sporulation initiation is becoming clear, and a summary of the current speculative regulatory pathways are shown in Figure 2.9.

The decision to sporulate begins with a two-stage signal transduction mechanism. In the first stage, a transmitter protein, which in some way recognizes nutrient depletion, passes an activation signal to a receiver protein. At this time, the identity of the transmitter and the mechanism by which the transmitter senses the environment have not been substantiated, however, evidence indicates a potential role for GTP as the environmental stress signal [102, 77]. At any rate, when the differentiation signal is recognized, it is believed auto-




**Figure 2.9**: Regulatory pathway for sporulation initiation. Arrows indicate flow of the initiation signal. Circles indicate down-regulation. Adapted from Smith [100].

phosphorylation of the transmitter protein occurs. The phosphorylated transmitter transfers the activation signal to a receiver protein through a subsequent phosphorylation of the receiver protein. Currently, the identity of the receiver protein has been attributed to the *spo*0A gene product. Evidence pointing to Spo0A as the receiver protein includes the sequence homology exhibited between Spo0A and known signal receiver proteins in other bacterial systems [100], and to the fact that the Spo0A protein is recognized as a master regulator of late growth processes and early sporulation events [81].

Once the receiver has been activated, a combination of events ensues, triggering the cascade of morphological and biochemical events associated with differentiation. The receiver protein, Spo0A, is a DNA binding protein that interacts with the promoter of at least one gene, *abrB*. The gene product of *abrB* plays a major role in the sporulation cascade as a repressor of many sporulation related genes [100]. The AbrB protein is expressed during exponential growth and is able to bind to a set of similar promoter regions, rendering them untranscribable. The interaction of Spo0A with the promoter region of *abrB* interferes with the expression of AbrB and allows for transcription of the previously repressed sporulation genes [115].

A particularly important product, expressed only after inhibition of AbrB, is the sigma factor, $\sigma^H$. This product of the *spo0H* gene is a protein that in combination with the "core" RNA polymerase enzyme, is able to transcribe a specific set of sporulation-associated genes. The altered selectivity of the RNA polymerase holoenzyme complex, E$\sigma^H$, is due to a selective association between the sigma factor and the promoter region of the differentiation genes. Utilization of different sigma factors is a major regulatory mechanism in the response of *Bacillus* to adverse environmental conditions [18]. In the case of $\sigma^H$, mutational analysis has shown that this sigma factor must be functional for sporulation initiation, due to the fact that the spore development pathway depends on products whose transcription is directed by the E$\sigma^H$ complex [62, 102]. The cascade of sporulation related events continues with the sequential expression of products involved in formation of the endospore, along with additional sigma factors, needed to direct sporulation.

Although the regulatory mechanism for control of sporulation initiation is becoming clear, it is difficult to translate the detailed regulatory information into a useful mechanistic model due to the lack of kinetic understanding for the various developmental steps. As was mentioned previously, some researchers are utilizing highly-structured models to describe sporulation [104, 56]. However, the large number of parameters, and the lack of measurable states brings into question the feasibility of model identification for these single cell models. Prior to the single-cell models, work concerning sporulation modelling has depended on empirical approaches to describe sporulation initiation.

Researchers have shown that several *Bacillus* species form spores during exponential growth in a chemostat [21, 92]. It has also been shown that sporulation initiation decreases with increased growth rate. The observation that spore production did occur at steady state in an exponentially-growing continuous culture contradicted the commonly held belief that sporulation was a threshold event, in which vegetative cells committed to spore formation in mass when the culture environment could no longer support a basal growth rate. In light of the knowledge concerning the sporulation regulatory mechanism, it seems likely that the inverse relationship between sporulation and growth rate at steady state is a result of interactions between the nutrient-limitation signal molecule (GTP) and the nutrient-sensing transmitter protein. This can be described using an equilibrium relationship:

$$SP \rightleftharpoons S + P^* \rightarrow \text{sporulation} \tag{2.8}$$

where $SP$ represents the inactive nutrient-protein complex and $P^*$ is the activated transmitter protein initiating sporulation. Lower growth rate implies lower nutrient concentration, which would shift the equilibrium relationship toward activated transmitter. The belief that mass sporulation was attributed to a threshold event could be of a sudden shift in the equilibrium relationship due to rapid exhaustion of limiting substrate in batch culture. This theory is consistent with the low, but non-zero sporulation rate observed in high dilution rate continuous cultures, as well as the mass sporulation seen at low dilution rate and in batch growth.

In an attempt to model the apparent dependence of sporulation on growth rate, Dawes and Thornley [22] defined the specific rate of sporulation to represent the rate of loss of vegetative cells, $X_v$ due to sporulation:

$$\mu_s = -\frac{1}{X_v}\frac{dX_v}{dt} \tag{2.9}$$

Based on observation of steady-state sporulation in *B. subtilis*, they postulated a linear relationship between the vegetative growth rate, $\mu_v$, and $\mu_s$:

$$\mu_s = \alpha_s - \beta_s \mu_v \tag{2.10}$$

The parameter $\alpha_s$ represents the maximum rate of sporulation and $\beta_s$ modulates the inhibitory effect of vegetative growth on the rate of sporulation. This relationship shows the decrease in sporulation observed when growth rate is high. The dependence of $\mu_s$ on $\mu_v$ provides an implicit dependence of sporulation on limiting substrate concentration through Equation 2.2.

Dawes and Thornley were able to accurately represent steady-state sporulation data for *B. subtilis* using the linear relationship in Equation 2.10. However, batch experimental data in combination with dilution-rate step tests, indicate that the linear relationship does not hold when limiting substrate concentration reaches zero. Therefore, instead of assuming a linear relationship between growth and sporulation, an experimentally-fit polynomial expression has been utilized to model the dependence of sporulation commitment, $\mu_s$ on limiting substrate concentration. In a general formulation, the empirical sporulation function can be represented by:

$$\mu_s = \sum_{j=0}^{n} \gamma_j \phi_j(s_i) \tag{2.11}$$

where $\phi_j$ represents the $j$th basis function and $\gamma_j$ is the experimentally determined basis function weight. Utilization of a general polynomial approximation has proved effective in describing sporulation commitment during both steady state and transient growth, and avoids the assumption of linear dependence of sporulation on growth rate, which was made by Dawes and Thornley. A detailed discussion of the estimation of the sporulation function is given in Section 4.3.2.

**Spore Maturation**. Along with the description of the sporangium birth through sporulation initiation, an understanding of the mechanisms by which cells leave the population is required to complete the population description. For this study, it is assumed that there is no significant loss of cells resulting from death of sporangia. There is, however, a mechanism by which sporangia leave the population. At the conclusion of the sporulation process, cells have been completely transformed from the vegetative state to dormant spores. Therefore, by definition, the maturation time represents a division between the sporangium population and the unstructured spore state. Even though maturation does not represent the "death" of a sporangium, it is a mechanism by which sporangia leave the population, and therefore, it is convenient to treat it as a death term in the population balance equation.

After the completion of the differentiation process, sporangium cells are considered to be mature spores, capable of renewing the vegetative growth process through germination. While the point where a sporangium cell has truly reached maturity is somewhat arbitrary, for the purposes of this research it is defined to be the stage of development where sporangia acquire heat resistance. Based on the assumption that spore development occurs in a fixed amount of time, the age of spore maturation is defined as $\tau_m$. The period required for *B. subtilis* to achieve maturity has been estimated by several researchers to be in the range of 8–10 hours [76, 63]. Recognizing that the true time of spore maturation is vague, and to account for any species related differences between *B. licheniformis* and *B. subtilis*, $\tau_m$ has been treated as an adjustable parameter. It is also assumed that the maturation time is relatively insensitive to changes in culture conditions.

The rates of sporulation commitment and spore maturation are tied into the population balance equation by definition of the birth and death terms in

Equation 2.7.

$$\frac{\partial X_s}{\partial t} + \frac{\partial X_s}{\partial \tau} = \delta(\tau)\mu_s X_v - \delta(\tau - \tau_m) X_s \tag{2.12}$$

The Dirac delta functions associated with birth and death account for the fact that the processes of sporulation commitment and maturation act as a point source and sink, respectively.

**Antibiotic Formation Model.** The observation that secondary metabolites are most commonly found in organisms that utilize sporulation mechanisms has led to considerable speculation about the role of secondary metabolites in the process of cellular differentiation [70, 57, 10]. The role played by secondary metabolism in the differentiation process is still a matter of controversy, and although the sporulation regulatory mechanism is becoming clear, the exact connection of the late growth products to sporulation is still unknown [102].

Initially, speculation that antibiotic compounds were essential to sporulation was based on several experimental observations. First, it was noted that the production of most peptide antibiotics, including bacitracin, coincided with sporulation initiation [9, 10]. Growth studies examining the connection of spore formation to antibiotic production noted that catabolite repression affected both antibiotic expression and sporulation similarly [65, 47, 9]. Majumdar and Bose [66] were able to show simultaneous derepression of production of the peptide mycolinic acid and sporulation in *B. subtilis* using cyclic dibutryl GMP. Further, genetic studies showed mutants that were blocked early in the sporulation sequence were also antibiotic negative [91]. These observations from genetic and biochemical experiments lead Hodgson [52] to speculate that peptide antibiotics were required for normal sporulation.

Although much evidence points to a connection between sporulation and antibiotics, Katz and Demain [57] conclude that none of the evidence discussed above conclusively prove that the antibiotics are required for successful spore formation. Nakano and Zuber [75] point out that the evidence that Hodgson's speculation is based on could be explained by the presence of parallel regulatory pathways for spore formation and antibiotic expression starting at a common origin. This scenario would explain spo$^-$ and antibiotic$^-$ phenotypes of early blocked mutants, and the production of antibiotics in sporulation late-blocked mutants. Schaeffer [91] explains that a causal relationship between antibiotics and spore formation is difficult to prove, but easy to disprove. To disprove the necessity of a particular compound in sporulation, a mutant must be isolated whose phenotype is antibiotic$^-$, and yet, is able to sporulate normally. Antibiotic-negative mutants that sporulate normally have been discovered in several cases [84, 67]. In particular for bacitracin, Haavik and Thomassen [46] were able to isolate a sporulating mutant that was bacitracin deficient.

Although it is likely that bacitracin is not essential to sporulation in *B. licheniformis*, the evidence indicates that the regulatory pathways for spore formation and antibiotic production overlap in some fashion [57]. Before attempting to describe a model for antibiotic formation, it is helpful to access the experiences of previous researcher involved in bacitracin production. Reviews of the early fermentation studies are provided by Hickey [51] and Weinberg [110]. A more recent review of bacitracin production is given by Frøyshov [35].

In one of the first studies of fermentation conditions promoting bacitracin production, Hendlin [50] studied growth and antibiotic production of a strain of *B. subtilis* in shake flasks using various synthetic media. He reported simultaneous growth and antibiotic production. No attempt was made to analyze any connection between sporulation and secondary metabolism. However, the effect of aeration on bacitracin production was examined by varying medium volume in shake flasks. Hendlin observed increased rate of bacitracin production in the low-volume flasks, and based on these results he concluded increased aeration improved production.

Bernlohr and Novelli [9] studied bacitracin production during batch growth on minimal medium in shake flasks. They noted that bacitracin production was inhibited by high concentrations of glucose ($\approx 40$ g/l), and they attributed the inhibition to lowering of pH due to organic acid production. The level of inhibition was related to the agitation rate of the shake flasks. Fast-shake agitation lead to significant inhibition of bacitracin production, slow-shake agitation and stagnant conditions allowed for production of bacitracin after approximately 48 hours. By limiting the glucose concentration ($\approx 5$ g/l), Bernlohr and Novelli were able to stimulate bacitracin production at approximately 10 hours, and the dependence of antibiotic production on aeration rate was eliminated. These researchers observed that bacitracin was only produced in cultural conditions that eventually supported spore formation. Their explanation was that bacitracin regulation was connected to spore formation. However, Bernlohr and Novelli note that they did not find a direct proportionality between the quantity of spores produced and the bacitracin titer.

Haavik [43] observed bacitracin production concomitant with growth in a synthetic medium that lacked glucose. In similar experiments with glucose added to the medium, bacitracin production was retarded, appearing after a delay proportional to the concentration of glucose added. However, the maximum bacitracin titer was positively affected by the addition of glucose, and this was attributed to reduction of the high pH observed during the late growth in medium without glucose. Haavik suggests that bacitracin is not inhibited by glucose, but rather, the low pH induced by organic acid production resulting from glucose metabolism. To test this theory, Haavik used $CaCO_3$ buffered growth medium to show that the repression effect could be eliminated in the presence of glucose.

After more experimental work, Haavik [44] modified his theory of pH dependent inhibition of bacitracin production. It was discovered that pH alone was not responsible for reduced bacitracin titers. Cultures acidified with inorganic acids did not exhibit inhibition of bacitracin production. The repression effect was only observed when pyruvate and acetate where used for acidification. Additionally, the inhibitory effect of these organic acids, which result from metabolism of glucose, was only observed when the pH of the culture was allowed to drop. Haavik concluded that repression of bacitracin results when undissociated organic acids penetrate the cell, creating a lower internal cell pH. The conclusion that it is possible to produce bacitracin during growth is also asserted, and that previous studies have not seen this phenomenon because of repression due undissociated organic acids. However, it is important to note that Haavik did not examine the relationship between organic acid addition and sporulation. Therefore, it is difficult to rule out the possibility that sporulation was occurring at the same time as growth, resulting in bacitracin production.

Flickinger and Perlman [32] attempted to analyze the effects of oxygen-enriched aeration on the production of bacitracin by *B. licheniformis* in complex media. Oxygen enrichment had wide-ranging effects on growth and antibiotic production. The maximum growth rate declined and sporulated sooner than in normal air-sparged control cultures. Also observed was a reduced rate of carbohydrate utilization as oxygen transport was improved. In contrast to the reduced growth rate, increased aeration increased the final antibiotic yield, and significantly increased the rate of bacitracin synthesis. Flickinger and Perlman suggest that the oxygen enrichment simulates conditions of slow carbohydrate feeding, and thus, reduce any catabolite repression of antibiotic production created during growth on carbohydrates.

The question of whether bacitracin production was possible during active growth lead Hanlon and Hodges [47] to look into the relationship between bacitracin production and growth rate. These researchers grew *B. licheniformis* on a minimal-salts medium, while altering the defined carbon, or nitrogen source. Five carbon sources were analyzed, ranging from glucose, which was an excellent growth source, to lactate, which promoted growth at about 1/4 the rate of glucose. It was observed that the carbon sources that allowed rapid growth did not promote bacitracin synthesis, and slow growth on poor carbon sources yielded bacitracin production. The propensity to produce bacitracin during exponential growth was correlated with the maximal growth rate supported by the carbon source. It was also observed that the quantity of spores produced corresponded to the growth rate in the same manner as bacitracin production. Based on these observations, Hanlon and Hodges claim it is obvious that bacitracin can be produced during exponential growth. However, they are careful to point out that this does not mean that bacitracin production is unassociated with sporulation.

From the review of experimental studies of bacitracin production, it is clear that the regulatory mechanisms for bacitracin production and sporulation are not completely coupled. Bernlohr and Novelli [9] indicate that cultural conditions requiring sporulation were required for bacitracin expression, however they were not able to directly correlate the bacitracin titer to the extent of spore production. Haavik [43, 44] claims bacitracin can be produced during exponential growth, implying that differentiation is not required for antibiotic production, even though the rate of sporulation was not monitored. The results of Hendlin [50] and Flickinger and Perlman [32] suggest a connection between the increased aeration and increased bacitracin production. This observation conflicts with the observations of Bernlohr and Novelli [9], who claim increased mixing, implying improved oxygen transfer, leads to inhibition of antibiotic production.

The conflicting experimental results indicate a complex regulatory mechanism for bacitracin production. In spite of the assertions of Haavik [43] that bacitracin production is not necessarily sporulation dependent, the remaining experimental evidence indicating a link between differentiation and antibiotic production are impossible to ignore. The consensus seems to be that antibiotics and other *Bacillus* secondary metabolites are coupled to the differentiation mechanism at the point of initiation. Smith [100] proposes that expression of secondary metabolites that are associated with sporulation, but are not essential to the differentiation process, is controlled by a regulatory pathway that is distinct from sporulation regulation, but shares a common origin. A possible regulatory structure is shown in Figure 2.10. A branched regulatory mechanism could explain the seeming dependence of bacitracin on sporulation, while still allowing for independent regulation of antibiotic production.

For fermentation experiments carried out in this study, detailed in Chapter 4, a dependence of bacitracin production on aeration was observed, indicating some effect of dissolved oxygen concentration on bacitracin synthesis. Two aeration levels were utilized during bacitracin fermentations. At high aeration, growth and sporulation were observed, however, bacitracin synthesis was highly repressed, with no measurable bacitracin produced. Utilizing the




**Figure 2.10**: Possible connection between the regulatory pathways for sporulation and antibiotic production.

same growth medium with reduced air flowrate and slower agitation speed, growth and differentiation were observed, along with production of bacitracin shortly after the onset of stationary phase. The observation of repressed antibiotic synthesis with increased aeration agrees with results obtained by Bernlohr and Novelli [9] and conflicts with the observations of Flickinger and Perlman [32] and Hendlin [50]. Possible explanations for the discrepancy could be differences attributable to strain variations, media composition, or shear sensitivity.

In light of the fact that the sporulation pattern and the connection of antibiotic production to that pattern are not well understood, an empirical model for the rate of bacitracin synthesis is appropriate. To capture the dependence of bacitracin production on sporangium cell age, a polynomial expression in sporangium age has been used to describe the specific bacitracin formation rate:

$$\mu_a = \sum_{j=0}^{m} \alpha_j L_j(\tau) \tag{2.13}$$

The antibiotic production coefficients, $\alpha_1, \cdots, \alpha_m$, are experimentally fit, allowing for an empirical age-based idiolite formation expression. This approach is similar to that of Shu [96], except that the age-dependence of product formation is isolated in the sporangium population, rather than the entire cell population.

For the fermentation experiments in this study, the dissolved oxygen concentration was not measured. Therefore, it is difficult to account for the dependence of bacitracin production on dissolved oxygen in the age-based model. Since dissolved oxygen is not considered, the bacitracin production parameters are based on the aeration operating conditions used for their estimation.

### 2.2.9 Segregated Fermentation Model

**Dynamic Equations**. The dynamic segregated fermentation model is developed by coupling the kinetic expressions for the various rate processes to the fluid dynamics model for the fermentation apparatus of interest. In this study, a stirred tank fermentation reactor has been utilized, and in the development of the segregated model equations the following assumptions have been made:

1. The fermentation broth is a dilute solution of constant density.
2. The stirred tank fermentation vessel is perfectly mixed.
3. The fermentor is maintained at constant temperature.
4. All rate process are limited by a single substrate.
5. Feed to the fermentor is sterile, and is fed at a flowrate $F_i$, with a limiting substrate concentration, $s_f$.




**Figure 2.11**: Continuous stirred tank fermentor with a material-balance control volume defined by the dotted line.

Based on a continuous stirred tank fermentor of liquid volume $V_r$, with a control volume as indicated in Figure 2.11, the modelling equations are formed by performing material balances on spore and vegetative cell concentrations, limiting substrate concentration, antibiotic concentration, fermentor volume, and sporangium cell number density. The resulting coupled integro-differential equations compromise the dynamic segregated fermentor model. With ˆ over a variable representing a dimensionless state, the model is presented in dimensionless form:

Spore Biomass

$$\frac{d(\hat{X}_e \hat{V}_r)}{d\hat{t}} = \delta(\hat{\tau}-1)\hat{X}_s(\hat{t},\hat{\tau})\hat{V}_r - \mu_g \hat{X}_e \hat{V}_r - \frac{F_o}{V_o}\hat{X}_e \quad (2.14)$$

Vegetative Biomass

$$\frac{d(\hat{X}_v \hat{V}_r)}{d\hat{t}} = (\hat{\mu}_v - \hat{\mu}_s)\hat{X}_v \hat{V}_r + \hat{\mu}_g \hat{X}_e \hat{V}_r - \frac{F_o}{V_{r0}}\hat{X}_v \quad (2.15)$$

Bacitracin

$$\frac{d(\hat{A}\hat{V}_r)}{d\hat{t}} = \int_0^1 \hat{\mu_a}\hat{X}_s\hat{V}_r\, d\hat{\tau} - \frac{F_o}{V_{r0}}\hat{A} \tag{2.16}$$

External Substrate

$$\frac{d(\hat{s}_e\hat{V}_r)}{d\hat{t}} = \frac{F_i}{V_{r0}}\hat{s}_f - \hat{\mu}_i\hat{X}_v\hat{V}_r - \frac{F_o}{V_{r0}}\hat{s}_e \tag{2.17}$$

Internal Substrate

$$\begin{aligned}\frac{d(\hat{s}_i\hat{V}_r)}{d\hat{t}} &= \hat{\mu}_i\hat{X}_v\hat{V}_r - \frac{y_v}{y_g}\hat{\mu}_g\hat{X}_e\hat{V}_r - \left(\hat{\mu}_v + \frac{y_v}{y_s}\hat{\mu}_s\right)\hat{X}_v\hat{V}_r \\ &\quad - \left(\frac{y_v}{y_a}\int_0^1 \hat{\mu}_a\hat{X}_s\hat{V}_r\, d\hat{\tau}\right) - \frac{F_o}{V_{r0}}\hat{s}_i \end{aligned} \tag{2.18}$$

Culture Volume

$$\frac{d\hat{V}_r}{dt} = \frac{F_i}{V_{r0}} - \frac{F_o}{V_{r0}} \tag{2.19}$$

Sporangium Number

$$\begin{aligned}\frac{\partial(\hat{X}_s\hat{V}_r)}{\partial\hat{t}} &= -\frac{\partial\hat{X}_s}{\partial\hat{\tau}}\hat{V}_r + \delta(\hat{\tau})\hat{\mu_s}\hat{X}_v\hat{V}_r \\ &\quad -\delta(\hat{\tau} - \hat{\tau}_m)\hat{X}_s\hat{V}_r - \frac{F_o}{V_{r0}}\hat{X}_s \end{aligned} \tag{2.20}$$

Definitions of the variables are provided in the nomenclature section. For convenience, the state and independent variables, with their dimensionless definitions, are summarized in Table 2.2. The specific kinetic rate functions are described in Table 2.3. For convenience, the dimensionless notation, $\hat{}$ will be dropped, and all equations are presented in dimensionless form unless otherwise noted.

Equations 2.14–2.19 represent the dynamic ordinary differential equations (ODE) modelling the non-segregated vegetative and endospore cell populations along with the substrate, culture volume, and antibiotic material balances. Equation 2.20 is a balance on the sporangium cell number, resulting in a hyperbolic partial differential equation representing the change in number of sporangium cells of age, $\tau$, at time $t$. The sporangium cell balance is

| Variable | Dimensionless Definition | Description |
|---|---|---|
| $t$ | $t/\tau_m$ | Time (hr) |
| $\tau$ | $\tau/\tau_m$ | Sporangium age (hr) |
| $X_e$ | $\frac{X_e}{y_v s_{e0}}$ | Endospore concentration (1/l) |
| $X_v$ | $\frac{X_V}{y_v s_{e0}}$ | Vegetative cell concentration (1/l) |
| $A$ | $\frac{A}{s_{e0}}$ | Antibiotic concentration (g/l) |
| $s_e$ | $\frac{s_e}{s_{e0}}$ | External substrate concentration (g/l) |
| $s_i$ | $\frac{s_i}{s_{e0}}$ | Internal substrate concentration (g/l) |
| $X_s(t,\tau)$ | $\frac{X_s \tau_m}{y_v s_{e0}}$ | Sporangium number density $(1/(\mathrm{l} \cdot \mathrm{hr}))$ |
| $V_r$ | $\frac{V_r}{V_{r0}}$ | Working reactor volume (l) |

**Table 2.2**: Definitions of the state and independent variables for the age-segregated antibiotic production model.

| Function | Description |
|---|---|
| $\mu_a$ | Antibiotic production rate (1/hr) |
| $\mu_g$ | Endospore germination rate (1/hr) |
| $\mu_i$ | Substrate internalization rate (1/hr) |
| $\mu_s$ | Sporulation rate (1/hr) |
| $\mu_v$ | Vegetative growth rate (1/hr) |

**Table 2.3**: Specific rate processes for the age-segregated antibiotic production model.

coupled to the vegetative cell balance through sporulation initiation, which is treated as a point birth source in Equation 2.20. Maturation of sporangia couples the sporangium population balance to the endospore balance, appearing as the input term, $X_s(t,1)$, in Equation 2.14. Finally, integral terms in Equations 2.16 and 2.18 represent the cumulative age-dependent antibiotic formation and substrate consumption rates, coupling the population balance to the antibiotic and substrate balances.

Defining $\mathbf{x} = [X_eV_r, X_vV_r, AV_r, s_eV_r, s_iV_r, V_r, X_sV_r]^T$, $\mathbf{u} = [F_i, F_o, s_f]^T$, and $\theta$ as a vector of model parameters, along with specification initial conditions allows the segregated modelling problem to be conveniently represented by:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \frac{\partial \mathbf{x}}{\partial \tau}, \mathbf{u}; \theta) \quad (2.21)$$

$$\mathbf{x}(0,\tau) = \mathbf{x}_0(\tau) \quad (2.22)$$

Solution of these model equations constitutes the segregated model predictions of the dynamic growth, differentiation and antibiotic production of *B. licheniformis* in perfectly mixed stirred tank fermentor.

**Steady-State Equations**. Steady-state operation of a continuous stirred tank fermentor involves continuous feed of nutrient medium, accompanied by continuous removal of fermentation broth. Continuous operation of a fermentor, known as chemostat operating mode [4], is a useful technique for accessing organism behavior in constant environmental conditions. In this study, steady-state experiments have been utilized to gain information for estimation of model parameters. Therefore, a steady-state formulation of the segregated model is required to predict the steady-state behavior of the *B. licheniformis* system.

The steady-state behavior of the segregated fermentation model is represented by the fixed points of the dynamic equations. For a stable system, it is possible to find fixed-point solutions by integrating the dynamic model from a known initial point in time until all dynamics have damped out. A more direct approach is to form the steady-state model by substituting zero for all time dependent terms in the dynamic model. Since the reactor volume is constant, along with the flow into and out of the fermentor, it is convenient to define the operating parameter, $D = F/V_r$, known as the dilution rate. Substitution of $\mathbf{0}$ for the time dependent terms, and introduction of $D$ into the dynamic modelling equations gives the steady-state model:

$$
\begin{aligned}
0 &= \delta(\tau - 1)X_s(t,\tau) - \mu_g X_e - DX_e && (2.23)\\
0 &= (\mu_v - \mu_s)X_v + \mu_g X_e - DX_v && (2.24)\\
0 &= \int_0^1 \mu_a X_s \, d\tau - DA && (2.25)\\
0 &= D\,(s_f - s_e) - \mu_i(s_e)X_v && (2.26)\\
0 &= -\frac{1}{y_g}\mu_g X_e - \left(\mu_v + \frac{1}{y_s}\mu_s\right) X_v \\
&\quad - \left(\frac{1}{y_a}\int_0^1 \mu_a X_s \, d\tau\right) - Ds_i && (2.27)\\
0 &= -\frac{dX_s}{d\tau} + \delta(\tau)\mu_s X_v - \delta(\tau - \tau_m)X_s - DX_s && (2.28)
\end{aligned}
$$

Cast in vector form, the steady-state model is represented by:

$$\mathbf{0} = \mathbf{f}(\mathbf{x}, \frac{d\mathbf{x}}{d\tau}, \mathbf{u}; \theta) \qquad (2.29)$$

At steady-state, the fermentor conditions are time invariant, and the dynamic modelling equations are reduced from a set of coupled ordinary differential and partial differential equations to a set of algebraic equations coupled to a differential equation representing the population balance. Solution of these equations for a given set of operating conditions results in the steady-state bacitracin production model.

### 2.2.10 Solution of the Segregated Fermentation Model

**Dynamic Model Solution**. Complete analytical solutions of the resulting segregated dynamic modelling equations are generally not possible due to the nonlinear nature of the problem. Therefore, development of a numerical solution technique is necessary for solution of the fermentor model. Fortunately, efficient computer codes exist for the solution of nonlinear ODEs [12]. The population balance, which is a PDE, adds a degree of complexity to the problem. In fact, efficient solution of the PDE and the integral terms in Equations 2.16 and 2.18 is the crux of the numerical solution task.

While a variety of numerical techniques exist for solution of PDEs, the hyperbolic nature of the population balance leads to propagation of discontinuities along characteristic lines in the domain, creating difficulties when using standard finite-difference and method of weighted residuals techniques [60]. However, the linear structure of the age-population balance allows for a semi-analytical solution of Equation 2.20:

$$X_s(t,\tau)V_r(t) = X_s(t-\tau,0)V_r(t-\tau)\exp\left[-\int_{t-\tau}^{t}\frac{F_o(s)}{V_r(s)}\,ds\right] \tag{2.30}$$

The semi-analytical solution shows the propagation of the boundary condition down the characteristic lines defined by $t-\tau=c$. These characteristic lines of unit slope, represent the fact that time and age progress at an identical rate. An illustration of the age-time domain, and the relative roles of the initial and boundary conditions are shown in Figure 2.12.

To utilize the semi-analytical solution, $X_s(t,\tau)$ is approximated by a linear combination of polynomial basis functions in $\tau$ with time-varying coefficients:

$$X_s(t,\tau) \approx \sum_{i=0}^{n}\gamma_i(t)L_i(\tau) \tag{2.31}$$




**Figure 2.12**: Age-time characteristic lines for the age-segregated population balance. The PDE initial condition provides the initial data for characteristics in the shaded region where $\tau > t$, and the PDE boundary condition provides the initial data for characteristics where $t > \tau$.

The function can be represented equivalently, and more conveniently, in terms of its values $X_{s_i}(\tau_i)$ at $n$ points, $(\tau_i, \cdots, \tau_n)$, in the domain, known as collocation points [30]. The $L_i(\tau)$ polynomials are chosen such that they are orthogonal to each other on the problem domain, and the collocation points are chosen to be the $n+1$ roots of the $L_{n+1}(\tau)$ polynomial.

Expansion of the sporangium number density in orthogonal polynomials offers the advantage of an efficient means by which the substrate consumption and antibiotic production integrals, present in Equations 2.16 and 2.18, can be evaluated. The integral terms are approximated using an optimal Gaussian quadrature rule:

$$\int_a^b g(\tau)\, d\tau \approx \sum_{i=0}^{n} w_i g(\tau_i) \tag{2.32}$$

The $w_i$ are integration weights based on the type of orthogonal polynomials basis functions chosen, and $g(\tau_i)$ is the integrand evaluated at the $i$th collocation point. The quadrature rule is exact for any polynomial integrand of order $2n+1$ or less. This approach offers the advantages of high accuracy and low computational load over other quadrature techniques, such as Simpson's rule and the trapezoidal rule [109].

As mentioned previously, the hyperbolic nature of the population balance allows propagation of discontinuities down characteristic lines. The semi-analytical solution provides a convenient means by which the deformation of a point on the sporangium number density function can be tracked from its entry at the population boundary to its maturation. Since it is most convenient to follow a point as it travels through the age-time domain, the nature of the problem suggests an adaptive solution grid is needed to avoid storage of the boundary condition, and re-solution of the population balance for each point in a fixed grid. Additionally, since the characteristic lines are of constant slope, the structure of the quadrature grid is preserved during adaptation, and therefore, the optimal quadrature rule applies regardless of grid adaptation.

One problem encountered with the adaptive characteristics scheme is the fact that the grid moves away from $\tau = 0$, creating a poor approximation of the sporangium age distribution near $\tau = 0$. The approximation becomes progressively worse as the characteristics move farther away from the boundary. To improve the approximation an elemental grid is appropriate, allowing for addition of new elements to the quadrature grid as space becomes available. The adaptive elemental grid is illustrated in Figure 2.13. At the initial time,




**Figure 2.13**: The adaptive characteristics grid. Solid lines represent element boundaries, and dotted lines are internal collocation points. At $t = t_1$ the first new characteristic is initiated. At $t = t_2$ an entire new element has been added and the oldest element is deactivated.

the domain of interest is discretized by an elemental grid of the desired accuracy. As time increases, the grid moves away from the boundary. When time has increased a period that is equal to the distance from the elemental boundary to the next collocation point, a new characteristic is introduced, with an initial value determined by the boundary condition:

$$X_s(t, 0) = \mu_s X_v \tag{2.33}$$

Initiation of a new characteristic line is depicted at time $t_1$. As time progresses further, new characteristics are initiated as each collocation point crosses into the right half plane of the time-age domain. Adaption continues until an entire element has been added to the grid, illustrated at $t_2$. At the same time as the new element moves into the age domain, the oldest element has moved out of the region of interest. The oldest element becomes inactive and is dropped. By constantly repeating elemental addition and deactivation, the age domain of interest is accurately approximated without accumulating unneeded characteristics.

Substitution of the approximate integral terms into the dynamic differential equations for the lumped fermentor states, and coupling of the discretized population balance equations reduces the dynamic fermentation model to a set of nonlinear, ordinary differential equations coupled to the semi-analytical solution to the population balance. While it is currently not possible to solve general nonlinear differential equations analytically, a variety of numerical integration packages exist for solution of coupled nonlinear ODEs. Therefore, the dynamic fermentation model has been reduced to a numerically tractable form.

### 2.2.11 Steady-State Segregated Model Solution

Similar to the dynamic model solution, the nonlinear nature of the fermentation model precludes a general analytical solution for the steady-state case. However, at steady-state there is a simple semi-analytical solution to the sporangium age distribution equation. Integration of Equation 2.28 with respect to sporangium age and substitution of the steady-state sporulation initiation expression gives the semi-analytical, steady-state sporangium age density:

$$X_{s_{ss}}(\tau) = \mu_s(s_{i_{ss}}) X_{v_{ss}} \exp\left[-D\tau\right] \tag{2.34}$$

This equations reflects the fact that the sporangium age distribution falls off exponentially at steady state due to the increased probability that a cell will wash out of the fermentor as it ages.

As with the dynamic model, the analytical solution to the population balance is utilized by discretizing the age-dependent sporangium density, and expanding the steady-state number density as orthogonal polynomials in $\tau$. Representing the steady-state distribution in terms of its value at the collocation points reduces the state state sporangium distribution to a set of algebraic equations:

$$X_{s_i}(\tau_i) = \mu_s(s_{i_{ss}}) X_{v_{ss}} \exp\left[-D\tau_i\right] \tag{2.35}$$

With this representation of the sporangium density function, the antibiotic production and substrate consumption integrals can be approximated using Gaussian quadrature. This reduces the steady-state model to a set of nonlinear algebraic equations. Similar to the dynamic case, general analytical solutions to the steady-state model are not possible, however, a variety of algorithms exist to solve sets of nonlinear algebraic equations.

**Segregated Model Solution Code: SEGMOD**. To obtain solutions for the dynamic and steady-state segregated fermentation models, the adaptive grid algorithm discussed in Section 2.2.10 was implemented as a FORTRAN-77 subroutine. The solution routine, called SEGMOD, was constructed in modular fashion, allowing easy portage to a variety of driver applications requiring model solutions.

SEGMOD was designed to handle both the steady state or the dynamic model solution, based on the user's request. To solve both problems, the solution routine needs the capability to generate numerical solutions to nonlinear algebraic equations for the steady-state problem, and to numerically integrate nonlinear ordinary differential equations for the dynamic model. To reduce the burden of software development, two public domain routines were utilized to handle the bulk of the computations. The subroutine HYBRD1, from the MINPACK subroutine library [74], uses a modified Powell-hybrid method to find a zero of a system of nonlinear algebraic equations. HYBRD1 was utilized to find the steady-state solutions of the segregated fermentation model, through the solution to Equations 2.23–2.28. The dynamic model differential equations were integrated using the the FORTRAN subroutine DDASSL [82]. DDASSL is designed to solve first-order differential algebraic systems using a variable stepsize, variable order, backward differentiation algorithm.

Since the primary numerical tasks are handled by DDASSL and HYBRD1, the routine SEGMOD serves as a management routine for the model solution task. However, coupling the adaptive grid algorithm to the normally self-contained integration routine, DDASSL, requires tight coordination between the processes of grid adaptation and the integration. A general flowchart for the SEGMOD model solution routine is shown in Figure 2.14. The model solution begins by obtaining the initial conditions for the dynamic model, or an initial guess for the steady-state solution. Then the user determined fermentor operating conditions are loaded. At this point, depending whether a steady-state, or dynamic simulation is desired, the routine directs the initial conditions and operating parameters to the appropriate canned software. In the steady-state case, the program flow is straightforward, with HYBRD1 solving the steady-state algebraic equations, returning to the calling routine after convergence to the fix-point solution. Program flow in the dynamic simulations is more complex, due to coupling of the adaptive sporangium grid to the DDASSL integration software. DDASSL integrates dynamic problems in a step-wise manner. In the SEGMOD routine, DDASSL is put in intermediate step mode, forcing the integration to stop at each successful integration step. At the stopping point, the progress of the integration is checked. If the integration has progressed to a characteristics addition point or elemental falloff, as shown in Figure 2.13, the sporangium grid is adapted appropriately. After the grid is adapted, control is returned to the integrator for continued progression of the solution. This process continuous until the final integration time is reached, at which point, the model solution is returned to the calling




**Figure 2.14**: Flowchart for the segregated model solution subroutine SEGMOD.

routine.

It is important to access the ability of the numerical solution scheme to accurately represent the dynamics of the fermentation model. In the case of the segregated model, the accuracy of the solution hinges on the ability of the adaptive grid to track the sporangium age distribution. Since the solution scheme utilizes an adaptive elemental grid, it is important to determine what level of grid refinement is necessary to accurately model the population dynamics. However, since additional grid elements increase the size and computational overhead of the problem, it is imprudent to overload the solution grid with more elements than necessary.

For computational convenience, the adaptive grid has been designed such that the elements are equally spaced, and the polynomial basis functions are identical on each element. Therefore, the accuracy of the solution is controlled by varying the number of elements making up the grid. To access the accuracy of the solution scheme, the effect of grid refinement on solution to the population balance has been analyzed. Batch fermentations of *B. licheniformis* are typified by a burst in sporulation initiation when substrate levels fall to critical levels. Since sporulation bursts lead to the introduction of steep fronts in the sporangium age distribution, the solution of the batch fermentation model poses a tough test for the numerical solution scheme. A simple way to visualize the effect of grid refinement on the population balance solution is to examine the variation in the age-distribution at constant time, as the number of elements increases.

Figures 2.15–2.17 show the predicted batch sporangium number density versus cell age at a given time, with varying numbers of quadratic elements. The three plot sequence compares the solution at early, medium, and late times in the batch simulation. In each case, as the number of elements increases the sporangium age density curve converges and eventually the addition of elements has an insignificant effect on the approximate solution. The increased accuracy at the high-element approximations is evident in the early and medium-time plots, due to the existence of a peak in the distribution. With a ten-element grid, the basic shape of the curve has been captured, and additional elements add little to the description. In the late-time case, the lack of a peak allows for a accurate low-element approximation to the distribution. Since it is necessary that the solution be accurate throughout the simulation, the grid was set to ten quadratic elements, allowing for a good approximation at all times.

After testing of the model solution routine, and analysis of the required grid spacing, solutions of the segregated model are available for simulation of fermentor operation. A simulation of a batch fermentation with nominal model parameters is shown in Figures 2.18 and 2.19. The lumped parameter states show the typical batch behavior, with rapid increase in the vegetative cell biomass until limiting substrate is exhausted. At the point of substrate depletion, vegetative cell death and sporulation phenomenon become active. Bacitracin is produced during the mass sporulation, and spores appear after the sporangia begin to mature.

The sporangium number density is a distributed parameter state, and complete representation of $X_s(t, \tau)$ requires a surface plot. The surface plot shown in Figure 2.19 shows the sporangium age distribution as it deforms in time for the batch fermentation. At the initial condition, there are few sporangia, however, as the nutrient is depleted, sporulation initiation creates a burst of newly born sporangia. After a short period of mass sporulation, the rate of spore initiation decreases and the peak in sporangium moves through the age-time




**Figure 2.15**: The effect of grid refinement on the sporangium number density solution early in a batch fermentation.

domain as the sporangia develop into mature spores.

### 2.2.12 Segregated Model Parameters

To finalize the segregated model, the parameters describing fermentation kinetics, stoichiometry, and sporulation maturation time must be determined. For the kinetic processes the parameters include:

- Maximum germination rate and substrate affinity: $\alpha_g$ and $K_g$.
- Maximum substrate internalization rate and external substrate affinity: $\alpha_i$ and $K_i$.




**Figure 2.16**: The effect of grid refinement on the sporangium number density solution midway through a batch fermentation.

- Maximum vegetative growth rate and internal substrate affinity: $\alpha_v$ and $K_v$.
- Cell death rate, $m_v$.
- Sporulation initiation coefficients: $\gamma_i,\ i = 1, n$.
- Antibiotic production coefficients: $\alpha_j,\ j = 1, m$.

For the growth and differentiation processes, the parameters relate the process rates to the limiting substrate concentration. Substrate consumption is related to growth, differentiation, and bacitracin production through stoichiometric




**Figure 2.17**: The effect of grid refinement on the sporangium number density solution late in a batch fermentation.

constants, $y_v$, $y_g$, $y_s$, and $y_a$. The final parameter is the spore maturation time, $\tau_m$, marking the time from sporulation initiation to maturation.

The lack of relevant literature data for the *B. licheniformis* system precludes the possibility of determining all model parameters from existing models or data. Therefore, it is necessary to estimate the parameters from experimental data generated for this study. Fortunately, sensitivity analysis indicates the model is insensitive to some parameters, allowing those parameters to be fixed at reasonable values. Then parameter estimation can concentrate on estimating the high sensitivity parameters. These topic are discussed in detail in Chapter 4.




**Figure 2.18**: Dynamic profiles of the lumped parameter segregated fermentation model states.

## 2.3 Development of the Structured Comparison Model

A useful tool for evaluating a model is comparison to other modelling approaches. In the case of antibiotic production by *Bacillus* an age-structured model exists in the fermentation literature. This model was developed by Blanch and Rogers to describe production of gramicidin S by *B. brevis* [11]. This model has been selected as a model suitable for comparison to the structured/segregated model developed in this research. Since the Blanch and Rogers (BR) model lacks segregation but includes structure, it was decided it would provide a valid test of the advantages and disadvantages of the segre-